Vol. 94, Sect. 14, 1957

PROTOCHORDATA

COMPILED BY

D. B. CARLISLE, M.A., D.Phil.

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE TWO SHILLINGS

December, 1959

SEPARATE SECTIONS OF THE ZOOLOGICAL RECORD

In addition to the complete bound volume, the separate Sections may be obtained singly, bound in printed paper covers.

The following are the Sections and their prices, viz.:-

| | 8. | d. |
|--------------------------------------|----|----|
| 1. Comprehensive Zoology | 3 | 0 |
| 2. Protozoa | 10 | 0 |
| 3. Porifera | 2 | 0 |
| 4. Coelenterata | 4 | 0 |
| 5. Echinoderma | 4 | 0 |
| 6. Vermes | 10 | 0 |
| 7. Brachiopoda | 4 | 0 |
| 8. Bryozoa | 2 | 6 |
| 9. Mollusca | 10 | 0 |
| 10. Crustacea | 6 | 6 |
| 11. Trilobita | 5 | 0 |
| 12. Arachnida | 10 | 0 |
| *13. Insecta | 50 | 0 |
| 14. Protochordata | 2 | 0 |
| 15. Pisces | 8 | 6 |
| 16a. Amphibia | 6 | 0 |
| 16в. Reptilia | 7 | 6 |
| 17. Aves | 10 | 0 |
| 18. Mammalia | 10 | 0 |
| 19. List of New Genera and Subgenera | 4 | 0 |

^{*} Obtainable only from the Commonwealth Institute of Entomology, 56, Queen's Gate, London, S.W.7.

14. PROTOCHORDATA

INCLUDING

ENTEROPNEUSTA, PTEROBRANCHIA, POGONOPHORA AND PHORONIDEA

BY

D. B. CARLISLE, M.A., D.Phil.

Evidence has been accumulating in recent years that the Graptolita are closely related to the Pterobranchia and not to the Coelenterata, with which they have been traditionally associated. From 1960, therefore, papers on Graptolita will be recorded in this section. Up to and including 1959 they will be recorded in Section 4 Coelenterata, where they have always been placed.

I.—TITLES

The year of publication is omitted where it is the same as the volume year of the "Record" namely (1957).

Abrikov, C. C. On the new type of invertebrates—Pogonophora. Adv. mod. Biol. Moscow 44 2:232-240 7 figs. [Russian.]

Alvariño, A. (1). Estudio del zooplancton de Mediterraneo Oxidentale. Bol. Inst. exp. Oceanogr. No. 81.

Alvariño, A. (2). Zooplancton del Atlantico Iberico. Bol. Inst. esp. Oceanogr. No. 82.

Anderson, J. M. see Guthrie, M. J.

Arzt, T. Die Erforschungsgeschichte der Chorda dorsalis und die Entstehung des Chordaten-Begriffes im 19, Jahrhundert. Nova Acta Leop. N.S. 17, 121 1955: 357–409.

Barrett, J. H. see Bassindale, R.

Barrington, E. J. W. The distribution and significance of organically bound iodine in the ascidian *Ciona intestinalis* Linnaeus. J. mar. biol. Ass. U.K. 36: 1-16, 1 pl. 1 fig.

Barrington, E. J. W. & Franchi, L. L. Organic binding of iodine in the endostyle of *Ciona intestinalis*. Nature, Lond. 177 1956: 432.

© Zoological Society of London. Vol. 94 Bassindale, R, & Barrett, J. H. The Dale Fort marine fauna. Proc. Bristol Nat. Soc. 29: 227-328 3 figs.

Berg, W. E. Chemical analyses of anterior and posterior blastomeres of *Ciona intestinalis*. Biol. Bull. Woods Hole 113: 365–375 2 figs.

Björnberg, T. K. S. (1). The occurrence of Amphioxides pelagicus (Günther) in the Fernand de Noronha Island plankton. Bol. Inst. oceanogr. S. Paulo 5 1954:83–86 1 fig.

Björnberg, T. K. S. (2). Sôbre quatro Tornárias do Atlântico e do Mediterrâneo. Bol. Inst. oceanogr. S. Paulo 6 1-2 1955: 197-213 3 pls. [English summary.]

Björnberg, T. K. S. & Forneris, L. Resultados científicos do cruzerio do "Bacpendé" e do "Vega" à ilha de Trinidade. Copelata 1. Publ. avuls. Inst. oceanogr. S. Paulo. Oceanogr. Biol. No. 1 1955: 1–68 17 pls.

Brewin, B. I. Ascidians of New Zealand. Part X. Ascidians from North Auckland. Trans. roy. Soc. N.Z. 84: 577-580 3 figs.

Burdon-Jones, C. Hemichordata. Enteropneusta. Family: Ptychoderidae. Tornaria larvae. Fich. Ident. Zoopl. Sheet 70: 1-6. Ceccatty, M. P. de. Les activités sécrétices du système nerveux central de l'*Amphioxus*. C.R. Acad. Sci. Paris **244**: 2645–2647.

Colwin, A. L., Colwin, L. H. & Philpott, R. E. Electron microscope studies of early stages of sperm penetration in *Hydroides hexagonus* (Annelida) and *Saccoglossus kowalevekii* (Enteropneusta). J. biophys. biochem. Cytol. **3**: 489–502 4 pl.

Colwin, L. H. see Colwin, A. L.

Dalcq, A. M. La métachromasie in vivo au bleu de toluidine dans l'oeuf et l'embryon d'Ascidie. Bull. Soc. zool. Fr. 82: 296-316 10 figs.

Davis, C. C. The Marine and Fresh-water Plankton (Ann Arbor) 1955: i-ix, 1-562 681 figs.

Deckert, K. Acrania see Streseman pp. 467-8.

Deflandre, G. & Deflandre-Rigaud, M. Micrascidites manip. nov., selérites de Didemnidés (Ascidies, Tunicier) fossile du Lutétien du Bassin parisien et du Balcombien d'Australie. C.R. Soc. géol. Fr. 1956: 47-9 figs. 1-5.

Deflandre-Rigaud, M. see Deflandre G.

Diehl, M. Die Ökologie der Ascidie Styela coriacea in der Kieler Bucht. Kieler Meeresforsch. 13: 59–68 5 pls.

Ebara A. (1). Physiological studies on the heart of an Ascidian, *Polycitor mutabilis*, Oka V. Action of Acetylcholine. Zool. Mag. Tokyo **62** 1953: 34.

Ebara, A. (2). Physiological studies on the heart of an Ascidian, *Polycitor mutabilis* Oka VI. The action of adrenalin. Zool. Mag. Tokyo 62 1953: 41–43.

Ebara, A. (3). Physiological studies on the heart of an Ascidian, *Polycitor mutabilis* Oka VII. The action of nicotine. Zool. Mag. Tokyo 62 1953:163–164.

Ebara, A. (4). Some observations on the heart-beat of the compound Ascidian, *Perophora orientalis* Ärnbäck. Sci. Rep. Tokyo Kyoiku Daig. 8B 124-130: 75-85 2 figs.

Ebara, A. (5). The electrocardiagram of the compound Ascidian, *Perophora orientalis* Ärnbäck. Sci. Rep. Tokyo Kyoiku Diag 8B: 126–37 3 pls. 1 fig.

Forneris, L. (1). The geographical distribution of the Copelata. Ann. Acad. bras. Cienc. 29:273-284.

Forneris, L. (2). Phoronidea. Family Phoronidea. Actinotroch larvae. Fich. Ident. Zoopl. Sheet 69: 1-4.

Forneris, L. see Björnberg, T. K. S.

Franchi, L. L. see Barrington, E. J. W.

Franzén, A. On spermiogenesis, morphology of the spermatozoon, and biology of fertilization among invertebrates. Zool. Bidr. Uppsala 31:355-482 6 pls. 223 figs.

Gautier, Y. V. Bionomie du Banc de Magaud. Rec. Trav. Stat. mar. Endoume 21: 28-40.

Gideon, P. W., Menon, P. K. B., Rao, S. R. V. & Jose, K. V. On the marine fauna of Gulf of Kutch: A preliminary survey. J. Bombay nat. Hist. Soc. 54: 690-706 1 pl. 5 figs. map.

Glover, R. S. An ecological survey of the drift-net herring fishery off the north-east coast of Scotland. Bull. mar. Ecol. 5 (39): 1–43 8 pls. 6 figs.

Godeaux, J. Contribution à la connaisance de la glande neurale des Polyclinidae (Ascidies Aplousobranches). Ann. Soc. zool. Belg. 87:75–89 7 figs.

Goodbody, I. Nitrogen excretion in ascidiacea. I. Excretion of ammonia and total non-protein nitrogen. J. exp. Biol. 34:297-305.

Grosch, D. S. & Smith, Z. H. X-ray experiments with *Molgula manhattensis*: adult sensitivity and induced zygotic lethality. Biol. Bull. Woods Hole 112: 171–179.

Gruner, H.-E. Tunicata se Stresemann, pp. 462-6.

Guthrie, M. J. & Anderson, J. M. General Zoology. New York (John Wiley & Sons), pp. xv+708 1 chart text illust.

Hattori, S. see Nakai, Z.

Honjo, K. see Nakai, Z.

Ikomoto, N. see Kawaguti, S.

Ionkov, V. I. Impressions of the lower Sarmatian Ascidiae. Priroda 6: 97–98 2 figs. [Russian.]

Ivanova-Kasa, O. M. Alterations in the size of nuclei in the ontogenesis of an ascidian *Dendrodoa grossularia* (Bened). Trans. Soc. Nat. St. Pétersb. (Léningr.) 69 1947: 101–136 24 figs. [Russian with English summary.]

Ivanov, A. V. (1). Neue Pogonophora aus die nordwestlichen Teil des Stillen Ozeans. Zool. Jb. Syst. **85**: 431–500 59 figs.

Ivanov, A. V. (2). On the systematic position of Pogonophora. Syst. Zool. 5: 165–173.

Ivanov, A. V. (3). Materials on the embryonic development of Pogonophora. Zool. Zh. 36: 1127-1144. [Russian with English summary.]

Jägersten, G. On the larva of Siboglimum. Zool. Bidr. Uppsala 32: 67-69 3 pl. 1 fig.

Jose, K. V. see Gideon, P. W.

Kawagati, S. & Ikomoto, N. Electron microscopy of the heart muscle from the squirt, Ciona intestinalis. Biol. J. Okayama Univ. 4: 93-101.

Kidachi, T. see Nakai, Z.

Kirkegaard, J. B. (1). Pogonophora. Galathealinum bruunii n. gen. n. sp., a new representative of the class. Galathea Rep. 2: 79-83 2 figs.

Kirkegaard, J. B. (2). Pogonophora. First records from the eastern Pacific. Galathea Rep. 2: 183–6 4 figs.

Kluge, H. On the occurrence of Rhabdopleura in the Barents Sea. Trud. Murmansk. Biol. Sta. Acad. Nauk. 1 1948: 276–283. [Russian.]

Kobayashi, S. see Mawatari, S.

Kott, P. (1). The ascidians of Australia. II. Aplousobranchiata Lahille: Clavelinidae Forbes and Harvey and Polyclinidae. Aust. J. mar, freshw. Res. 8:64–110 31 figs.

Kott, P. (2). The sessile Tunicata. John Murray Exped. Sci. Rep. 10: 129–149.

Kudoh, S. see Nakai, Z.

Lefevere, S. & Leloup, E. Observations biologiques dans le Port d'Ostende. Mém. Inst. Sci. nat. Belg. No. 133 1956: 1-157 3 pls.

Leloup, E. see Lefevere, S.

Lönöy, N. A comparative anatomical study of *Phoronis ovalis* Wright from Norwegian, Swedish and Brazilian waters. Univ. Bergen Årb. naturv. R. 1953 2 1954: 1–29 29 figs.

Mancuso, V. La distribuzione del glicogeno lungo lo sviluppo dell'uovo di *Phallusia mamillata*. Ric. sci. 26 1956: 2756–1760 7 figs. [French, English and German summaries.]

Marine Biological Association of the U.K. Plymouth Marine Fauna, Edit. 3, Plymouth (Mar. Biol. Assoc.).

Markman, B. On the peripheral nervous system of Ascidians. Acta Zool. Stockh. 39 (1):13-18 6 figs.

Mawatari, S. & Kobayashi, S. Seasonal settlement of animal fouling organisms in Ago Bay, middle part of Japan I. Misc. Rep. Res. Inst. nat. Resour. Tokyo, No. 35 1954: 37-47 11 figs. No. 36 1954: 1-8 2 pls. 6 figs.

Meel, L. van see Lefevere, S.

Menon, P. K. B. see Gideon, P. W.

Millar, R. H. (1). Ascidians from Mozambique, East Africa. Ann. Mag. nat. Hist. (12) 9:913-932 15 figs.

Millar, R. H. (2). Further notes on ascidians from Sierra Leone. Ann. Mag. nat. Hist. (12) 10:369–377 5 figs.

Minganti, A. (1). Studio cariologico di ibridi di Ascidie. Ric. sci. 25 1955: 2625–2630. [French, English and German summaries.]

Minganti, A. (2). I cromosomi nei Tunicata. Boll. Zool. 23 1956: 299–315. Nakai, Z., Hattori, S., Honjo, K., Watanabe, T., Ohta, H., Kidachi, T., Okutani, T., Umeda, K. & Kudoh, S. A preliminary report on the biological survey in the "Kuroshio" area, south of Honshu, June-July, 1955. Rec. oceanogr. Wks Jap. Special No. 1:159-196 46 figs.

Nogusa, S. The chromosomes of the Japanese lancelet, *Branchiostoma* belcheri (Gray) with special reference to the sex-chromosomes. Annot. zool. jap. 30: 42–46 12 figs.

Ohta, H. see Nakai, Z.

Oka, H. (1). On a new species of *Polycitor* from Japan with some remarks on its mode of budding. Annot. zool. Jap. 21 1942:155–162.

Oka, H. (2). Metamorphosis of *Polycitor mutabilis* (Ascidiae compositae). Annot. zool. Jap. 22 1943: 54–58.

Oka, H. & Usui, M. On the growth and propagation of the colonies in *Polycitor mutabilis* (Ascidiae compositae). Sci. Rep. Tokyo Bunrika Diagaku Sec. B. No. 103 1944: 23–53.

Oka, H. & Watanabe, H. (1). Colony-specificity in compound Ascidians as tested by fusion experiments (a preliminary report). Proc. Japan Acad. 33: 657-659.

Oka, H. & Watanabe, H. (2). Vascular budding, a new type of budding in *Botryllus*. Biol. Bull. Woods Hole 112: 225-240 11 figs.

Oka, H. & Watanabe, H. (3). Bourgeonnement vasculaire, nouveau mode de reproduction asexuée chez de Botryllidés. C.R. Soc. Biol. Paris 150: 1841–1843.

Okutani, T. see Nakai, Z.

Ortolani, G. (1). Azione della tripsina sul cortex dell'uovo di Phallusia mamillata. Ric. sci. 27: 1175–1178. [Italian with French, English and German summaries.]

Ortolani, G. (2). Rimozione dell' autosterilità in *Ciona intestinalis* mediante trattamento con versene. Ric. sci. 27 (7): 2150–2153. [Italian with French, English and German summaries.]

Ortolani, G. (3). Sviluppo dell' uovo di Ascidie dopo trattamento con tripsina. Ric. sci. 27: 1868–1873 5 figs. [Italian with French, English and German summaries.]

Ortolani, G. (4). Esperimenti di merogonia nell'uovo fecondata di Ascidie. Ric. sci. 27: 2475–2478. [Italian with French, English and German summaries.]

Pérès, J. M. (1). Ascidies récoltées dans les parages des Baléares par le "Professeur Lecaze-Duthiers" (Première partie: Majorque et Minorque). Vie et Milieu Suppl. 6: 177-184 1 fig.

Pérès, J. M. (2). Ascidies récoltées dans les parages des Baléares par le "Professeur Lecaze-Duthiers" (Deuxième partie : Iviza et San Antonio). Vie et Milieu Suppl. 6 : 223-224 1 fig.

Pérès, J. M. & Picard, J. Recherches sur les peuplement benthique. Ann. Inst. océanogr. Paris 24: 213-292.

Peyer, B. Über bischer als Fährten gedeutete problematische Bildungen aus den ologozänen Fischschiefern des Sernfatales. Schweizer. Paläontol. Abh. 73: 5-34 6 pl. 19 text figs.

Philpott, D. E. see Colwin, A. L.

Rao, K. P. (1). Enteropneusta from Madras and the Gulf of Manaar, with a note of the probable course of distribution of *Ptychodera flava* Eschscholtz. Proc. Indian sci. Congr. sci. Congr. Ass. 44 1955: 301. [Abstract].

Rao, K. P. (2). Two species of Enteropneusta from off the coast of Madras. Proc. Indian Sci. Congress 42 3 1955: 301. [Abstract].

Rao, S. R. V. see Gideon, P. W.

Reverberi, G. La citocromoossidasi lungo lo sviluppo delle ascidie Risultati che conseguono alla sua inibizione con l'azide sodico. Pubbl. Staz. zool. Napoli 29: 186– 212 1 pl.

Sabbadin, A, (1). Effetti dell'estirpazione delle gemme sulla durata del ciclo vitale in *Botryllus schlosseri* (Pallas). Boll. Zool. 23 1956: 331–342.

Sabaddin, A. (2). Il ciclo biologico di Ciona intestinalis (L.), Molgula manhattensis (De Kay) e Styela plicata (Lesueur) nella Laguna Veneta. Arch. Oceanogr. Limnol. 11: 1-28.

Scott, F. M. Regeneration and differentiation in buds of *Amaroecium* treated with nitrogen mustard. J. exp. Zool. **135**: 557–585 33 figs.

Skalkin, V. A. A new species of Ascidia of the Pacific—Syncarpa longicaudata sp. n. (Fam. Styelidae). Zool. Zh. 36: 297–299. [Russian with English summary.]

Sourie, R. Étude écologique des plages de la côte sénégalaise aux environs de Dakar (Macrofauna). Ann. Écol. supér. Sci. Dakar 3: 3-110.

Stresemann, E. & others. Exkursionsfauna von Deutschland. Wirbellose I. Berlin (Volk & Wissen Volks. Verleg) pp. xxxii+488 467 figs.

Tenbaum, E. Polarisationsoptische Beiträge zur Kenntnis der Gervebe von *Branchiostoma lanceolatum* (P.). Z. Zellforsch. **42** 1955: 149–192 38 figs.

Tokioka, T. Two new appendicularians from the eastern Pacific, with notes on the morphology of *Fritillaria aequatorialis* and *Tectillaria fertilis*. Trans. Amer. micr. Soc. **76**: 359–365 4 figs.

Trason, W. B. Larval structure and development of the oozooid in the ascidian *Euherdmania claviformis*. J. Morph. **100**: 509–526 9 pls.

Trégouboff, G. (1). Rapport sur les travaux concernant le plancton Meditérranéen publiés entre Novembre 1952 et Novembre 1954. Rapp. Comm. int. Mer Médit. 13 1956: 65–100.

Trégouboff, G. (2). Classe des Thaliacés. See Trégouboff, G. & Rose, M. pp. 537–570 pl. 197–205.

Trégouboff, G. (3). Classe des ascidies Blainville. See Trégouboff, G. & Rose, M. pp. 571-4 pl. 206.

Trégouboff, G. (4). Embranchement des Cephalocordés. *See* Trégouboff, G. & Rose, M. pp. 575–9 pl. 207.

Trégouboff, G. (5). Classe des appendiculaires Bronn (Pérennicordés Balfour). See Trégouboff, G. & Rose, M. pp. 512–536 pl. 192–196.

Trégouboff, G. & Rose, M. Manuel de planctologie méditerranéenne. Paris (Centre national de la Recherche scientifique).

Umeda, K. see Nakai, Z.

Usui, M. see Oka, H.

Watanabe, H. see Oka, H.

Vicentiis, M. de (1). Attività ossidativa dell'uovo di *Phallusia mamillata* in presenze di ossido di carbonio e di azide. Boll. Zool. 23 1956 (1957): 701–708.

Vicentiis, M. de (2). Influenza del CO sulla sviluppo di *Phallusia mamillata*. Ric. sci. 26 (10): 3093–3096 4 figs. [French, English and German summaries.]

Watanabe, T. see Nakai, Z.

Webb, J. E. Cephalochordata. Sci. Rep. Murray Exped. 10:121– 128 2 figs.

Whitear, M. Some remarks on the ascidian affinities of vertebrates. Ann. Mag. nat. Hist. (12) 10:338–348.

II.—SUBJECT INDEX

Reference to "Titles" is by the name(s) of the Author(s).

EVOLUTION & GENETICS

Phylogeny. — of chordates: White-

BRACHIATA

POGONOPHORA

GENERAL

Collections. — IVANOV (1).

STRUCTURE

Adult. — Abrikov, Ivanov (1).

DEVELOPMENT

Embryology. —IVANOV (3), JÄGER-

EVOLUTION & GENETICS

Phylogeny.—IVANOV (1, 2).

6

PHORONIDEA

GENERAL

Fauna lists—English Channel: MARINE BIOLOGICAL ASSOCIATION OF THE U.K.

Keys.—to larvae: Forneris (2).

STRUCTURE

Comparative anatomy.—Lönöy.

EVOLUTION & GENETICS

Variation.—Lönöy.

ECOLOGY & HABITS

Populations studies.—Lönöy.

PTEROBRANCHIA

GENERAL

Textbooks. — Guthrie & Anderson p. 543.

DEVELOPMENT

Sperm.—Franzén.

EVOLUTION & GENETICS

Affinities.—WHITEAR.

DISTRIBUTION

ARCTIC.—KLUGE.

ENTEROPNEUSTA

GENERAL

Fauna lists. — English Channel: MARINE BIOLOGICAL ASSOCIATION OF THE U.K. S. Wales: BASSINDALE & BARRETT: Madras: RAO (1).

Keys.—Burdon-Jones.

Plankton. — Björnberg (2), Burdon-Jones.

STRUCTURE

Larva.—Björnberg (2).

DEVELOPMENT

Sperm.—Franzén.

Fertilization.—Colwin et al.

ECOLOGY & HABITS

Distribution.—Rao (1, 2). Dispersal.—Rao (1).

DISTRIBUTION

NORTH TEMPERATE. — Mediterranean: Tornaria: Björnberg (2).

TROPICAL. — Atlantic: Tornaria: BJÖRNBERG (2); Indian Ocean: RAO (1, 2).

TUNICATA

GENERAL

Textbooks. — Guthrie & Anderson pp. 545-9; Ascidiacea: Trégouboff (3); Thaliacea: Davis, Trégouboff (2); Larvacea: Davis, Trégouboff (5).

History. — chorda dorsalis : ARZT.

Collections. — Ascidiacea: Brewin, Kott (1, 2), Millar (1, 2), Pérès (1, 2); Larvacea: Björnberg & Forneris, Tokioka.

Fauna lists.—English Channel: MARINE BIOLOGICAL ASSOCIATION OF THE U.K.; Ostend, Belgium: LEFEVERE et al.; Germany: GRUNER; India: GIDEON et al.; Ascidiacea: S. Wales; BASSINDALE & BARRETT; Red Sea and Arabia: Kott (1).

Keys. — Ascidiacea: Kott (1); Larvacea: Björnberg & Forneris.

Plankton. — ALVARIÑO (1, 2), BJÖRNBERG & FORNERIS, DAVIS, FORNERIS (1), GLOVER, NAKAI et al., TRÉGOUBOFF (1-5).

STRUCTURE

Adults. — Ascidiacea: neural gland: GODEAUX.

Larva, post larva. — Ascidiacea : Kott (1), Trason.

Nervous system. — Ascidiacea: Markman.

Endocrines. — Ascidiacea: God-

Biochemistry. — Ascidiacea: BARRINGTON.

Laboratory technique. — Ascidiacea: Dalcq.

Cytology, histochemistry.—Ascidiacea: Dalcq, Godeaux, Minganti (2), Ivanova-Kasas, Mancuso, Scott.

Electron micrography. — Ascidiacea: heart muscle: Kawaguti & Ikomoto.

PHYSIOLOGY

Feeding mechanisms. — Ascidiacea: Barrington, Barrington & Franchi.

Circulation, heart. — Ascidiacea: Ebara (1-5).

Excretion.—Ascidiacea: Good-Body.

Hormones. — Ascidiacea: Bar-RINGTON & FRANCHI.

Growth. — Ascidiacea: Oka & Usui, Sabbadin (1).

Regeneration. — Ascidiacea: Barrington, Barrington & Franchi, Berg, Vincentiis (2).

Enzymes. — Ascidiacea: VIN-CENTIIS.

Phagocytosis. — Ascidiacea: God-

House building. — Ascidiacea: Oka & Usui.

Radiation effects. — Ascidiacea: Grosch & Smith.

REPRODUCTION & SEX

Asexual reproduction. — Ascidiacea: Oka (1), Oka & Watanabe (2, 3), Sabbadin (1), Scott.

Fertility, autosterility. — Ascidiacea: Grosch & Smith, Ortolani (2).

Ovoviviparity. — Ascidiacea: Trason.

Breeding season. — Ascidiacea: Brewin.

Breeding cycle. — Ascidiacea: Diehl.

DEVELOPMENT

Egg, oogenesis. — Ascidiacea : Trason.

Sperm.—Franzén.

Fertilization.—Ascidiacea: Dalcq. Grosch & Smith, Ortolani (1-3).

Cleavage.—Ascidiacea: BERG, DALCQ.

Embryology. — Ascidiacea: Dalcq, Ivanova-Kasas, Trason. Chemical embryology. — Ascidiacea: Berg, Mancuso, Ortolani (1-4), Reverberi, Vincentiis (1, 2).

Organogeny. — Ascidiacea : Scott.

Larval stages, — Ascidiacea : Trason,

Metamorphosis.—OKA (2), TRASON.

Growth.—Ascidiacea: Ivanova-Kasas, Sabbadin (2); colony fusion: Oka & Watanabe (1).

Teratology. — Ascidiacea: Sabbadin (3), Scott.

EVOLUTION & GENETICS

Phylogeny. — Barrington, whitear, Ascidiacea: Barrington & Franchi.

Experimental genetics. — Ascidiacea: Oka & Watanabe (1).

Chromosomes. — Ascidiacea: MIN-GANTI (2).

Radiation effects. — Ascidiacea: Grosch & Smith.

ECOLOGY & HABITS

Ecology.—Ascidiacea: DIEHL, GAUTIER, SOURIE.

Environmental factors.—Ascidiacea: DIEHL.

Distribution. — Thaliacea: Alvariño (1, 2). Larvacea: Alvariño (2); temperature dependence: Forneris (1).

Growth. — Ascidiacea: DIEHL.

Annual eycle. — Ascidiacea: DIEHL, SABBADIN; Thaliacea, Larvacea: GLOVER.

Population studies. — Thaliacea: Alvariño (1, 2), Glover, Larvacea: Alvariño (2), Glover.

DISTRIBUTION

MARINE

NORTH TEMPERATE.—Mediterranean: Ascidiacea: Pérès (1, 2); Larvacea: ALVARIÑO (1, 2).

TROPICAL. — Atlantic: Ascidiacea: MILLAR (2); Larvacea: BJÖRNBERG & FORNERIS; East Pacific: Larvacea: Tokioka; Indian Ocean: Ascidiacea: Kott (2), MILLAR (1).

SOUTH TEMPERATE. — Ascidiacea: Brewin, Kott (1).

GEOLOGICAL

Pliocene. — Ascidiacea: Deflandre & Deflandre-Rigaud.

Oligocene. — Thaliacea: PEYER.

Lower Sarmatian of Russia.—Ascidiacea: Ionkov.

CEPHALOCHORDATA

GENERAL

Textbooks. — Davis, Guthrie & Anderson, Trégouboff (4).

History.—ARZT.

Fauna lists.—English Channel: MARINE BIOLOGICAL ASSOCIATION OF THE U.K.; S. Wales: BASSINDALE & BARRETT; Germany: DECKERT.

Collections.—Webb.

Plankton.—Björnberg (1), Trégouboff (4).

STRUCTURE

Adult.—Björnberg (1), Tenbaum, Webb.

Neurosecretion.—CECCATTY.

Nervous system.—CECCATTY.

Histology, Cytology.—CECCATTY, NOGUSA.

REPRODUCTION & SEX

Sex determination.—Nogusa.

Sperm, spermatogenesis. — Franzén, Nogusa.

EVOLUTION & GENETICS

Phylogeny.—Björnberg (1). Chromosomes.—Nogusa.

DISTRIBUTION

TROPICAL. — Atlantic: Björn-BERG (1).

III.—SYSTEMATIC INDEX

Reference to the "Titles" is by the name(s) of the Author(s).

Phylogeny of chordates.—Whit-EAR.

BRACHIATA POGONOPHORA

Systematic position.—Ivanov (2).

ATHECANEPHRIA

OLIGOBRACHIIDAE

fam. nov., Ivanov (1) p. 434: type gen. Oligobrachia.

Oligobrachia gen. nov., Ivanov (1) pp. 434, 439; type sp. O. dogieli; O. dogieli sp. nov. N.W. Pacific, Ivanov (1) pp. 434-9 figs. 1-5, embryology, Ivanov (3).

SIBOGLINIDAE

Siboglinum caulleryi sp. nov. N.W. Pacific, Ivanov (1) pp. 440-6 figs. 6-13, embryology, Ivanov (3); S. cinctutum sp. nov. N.W. Pacific, Ivanov (1) pp. 446-9 figs. 14-16; S. ekmani. Ivanov (1), embryology, Jägersten; S. fedotori sp. nov. N.W. Pacific, Ivanov (1) pp. 453-6 figs. 23-7; S. minutum sp. nov. N.W. Pacific, Ivanov (1) pp. 451-3 figs. 20-2; S. pellucidum sp. nov. N.W. Pacific, Ivanov (1) pp. 449-51 figs. 17-19; S. plumosum sp. nov. N.W. Pacific, Ivanov (1) pp. 456-61 figs. 28-30.

THECANEPHRIA

POLYBRACHIIDAE

Heptabrachia abyssicola, Ivanov (1); H. gracilis sp. nov. N.W. Pacific, Ivanov (1) pp. 461-5 figs. 31-4; H. subtilis sp. nov. N.W. Pacific, Ivanov (1) pp. 465-8 figs. 35-40.

Zenkevitchiana gen. nov., Ivanov (1) pp. 470, 475, type sp. Z. longissima; Z. longissima sp. nov. N.W. Pacific, Ivanov (1) pp. 470-5 figs. 41-5.

LAMELLISABELLIDAE

Lamellisabella johanssoni sp. nov. N.W. Pacific, Ivanov (1) pp. 475–483 figs. 46–52; L. zachsi, Ivanov (1).

SPIROBRACHIIDAE

Spirobrachia beklemischevi sp. nov. N.W. Pacific, Ivanov (1) pp. 483-9 figs. 53-8; S. grandis, Ivanov (1).

PHORONIDEA

PHORONIDAE

Actinotrocha branchiata=Phoronis mülleri; A. brownei, adult unknown; A. hippocrepia=P. hippocrepia; A. pallida=P. pallida: key and distribution, FORNERIS.

Phoronis ovalis, comparative anatomy, Lönöy.

PTEROBRANCHIA

RHABDOPLEURIDAE

Rhabdopleura annulata; R. compacta; R. grimaldi; R. manubialis; R. normanni; R. striata; R. sp. Harmer?=R. annulata; distribution and differences, Kluge.

CEPHALODISCIDAE

Cephalodiscus sperm, Franzén.

ENTEROPNEUSTA

PTYCHODERIDAE

Balanoglossus clavigerus, Björn-Berg (2), key to larva, distribution, Burdon-Jones,

Glossobalanus elongatus in Mediterranean and from Madras, RAO (2); G. marginatus distribution, Burdon-Jones; G. sarniensis distribution, Burdon-Jones, sperm structure, Franzén.

Ptychodera bahamensis ?=P. flava, Björnberg (2); P. flava ?=P. bahamensis, Björnberg (2), routes of dispersal, Rao (1).

Saccoglossus kowalevskii sperm penetration, Colwin et al.; S. madrasensis sp. nov. (nomen nudum) Madras, RAO (2) p. 301. Tornaria bournei key, distribution, Burdon-Jones; T. chierchiai I Brazil, first records from Atlantic, ?=T. morgani, Björnberg (2); T. dubia redescribed from Algiers, Björnberg (2); T. krohni redescribed from Algiers, ?=Balanoglossus clavigerus, Björnberg (2); T. meeki, T. mielcki key, distribution, Burdon-Jones; T. morgani, Björnberg (2); T. weldoni from Trinidad, Björnberg (2).

TUNICATA

Keys to German fauna, GRUNER.

ASCIDIACEA

CLAVELINIDAE

Clavelina baudiensis sp. nov. Australia, adult and larva, Kott (1) pp. 87-8 figs. 19-21; C. claviformis from Australia, larva figd., Kott (1); C. diminuta sp. nov. Australia, Kott (1) p. 89; C. enormis from Mozambique, larva figd., Millar (1), from Australia, larva figd., Kott (1); C. lepadiformis sperm, Franzén; C. sigillaria from N. Auckland, New Zealand, Brewin.

Distaplia australiensis, Kott (1); D. stylifera, Kott (1); D. viridis sp. nov. Australia, adult and larva, Kott (1) pp. 96-7 figs. 28-30.

Podoclavella australis, Kott (1); P. cylindrica from Australia, larva described, Kott (1); P. dagysa sp. nov. Australia, adult and larva, Kott (1) pp. 93-4 figs. 26-7; P. detorta from Arabia, Kott (2); P. meridionalis from Australia, larva, Kott (1); P. moluccensis from Australia, larva figd., Kott (1).

Stomozoa gen. nov., Kott (2) p. 131, type sp. S. murrayi; S. murrayi, sp. nov. Red Sea, Kott (2) pp. 131-4 figs. 2-8.

Stomozoinae subfam. nov., Kott (2) p. 131, type gen. Stomozoa.

Sycozoa cerebriformis; S. murrayi, larva, Kott (1); S. sigillinoides, Kott (1), from N. Auckland, New Zealand, Brewin; S. tenuicaulis, Kott (1).

POLYCITORIDAE

Archidistoma murrayi sp. nov. Arabia, adult and larva, Kott (2) pp. 134-5 fig. 9; A. parva from Australia, Kott (1); A. (Eudistoma) parva from Gulf of Aden, Kott (2).

Cystodytes dellechiaiae from Australia, Kott (1), Pérès (1).

Distaplia bermudensis from Mediterranean, Pérès (1).

Eudistoma angolanum from Australia, Kott (1), from Sierra Leone, larva, Millar (2); E. arenosum sp. nov. Australia, adult and larva, Kott (1) pp. 73-4 figs. 3-4; E. caeruleum from Mozambique, Millar (1); E. costai, Pérès (2); E. cyanea from Australia, Kott (1); E. elongata from Australia, larva figd., Kott (1); E. fantasiana sp. nov. Australia, adult and larva, Kott (1) p. 76 figs. 7-8; E. globosum sp. nov. Australia, adult and larva, Kott (1) pp. 72–3 figs. 1–2; E. mobiusi and E. passleroides from Mozambique, MILLAR (1); E. parvum from Australia, larva figd., Kott (1); E. plumbeum, Pérès (1); E. pyriforme from Australia, larva figd., KOTT (1); E. ramosum from Sierra Leone, adult and larva figd., MILLAR (2); E. rhodopyge from Mozambique, larva figd., MILLAR (1); E. renieri from Australia, larva figd., Kott (1); E. rubrum, PÉRÈS (1); E. tridentatum, Pérès (2).

Paradistoma cristallinum, Pérès (2).

Polycitor circes from Australia, KOTT (1); P. giganteum from Australia, larva figd., KOTT (1); P. gelatinosum sp. nov. Australia, larva figd., KOTT (1) p. 83 fig. 15; P. longitubis sp. nov. Australia, larva figd., KOTT (1) pp. 80-1 fig. 11; P. mutabilis sp. nov. Japan, OKA (1) pp. 156-9 figs. 1-9, budding and growth, OKA (1), OKA & USUI, metamorphosis, OKA (2), heart physiology, EBARA (1-3); P. subarborensis sp. nov. Australia, KOTT (1) pp. 81-2 fig. 14; P. translucida sp. nov. Australia, KOTT (1) p. 81 fig. 13.

Polycitorella mariae from N. Auckland, New Zealand. Brewin.

Sigillina maghalensis from Australia, Kott (1).

POLYCLINIDAE

Amaroucium areolatum, Pérès (1); A. brementi, Pérès (2); A. circumvollatum from N. Auckland, New Zealand, Brewin; A. constellatum regeneration, Scott; A. cottrelli sp. nov. N. Auckland, New Zealand, Brewin p. 578, fig. 1; A. nottii, A. phortax and A. thomsoni from N. Auckland, Brewin; A. turbinatum ecology, Gautier.

Aplidium altarium from Mozambique, larva figd., MILLAR (1); A. caeruleum var. argelense, PÉRÈS (1); A. lubricum from Mozambique, larva figd., MILLAR (1); A. petrense from Mozambique, A. savignyi var translucidum var. nov. Arabia; Kuria Muria Islands, Kott (2) pp. 142-3; A. violaceum from Arabia, larva figd., Kott (2).

Euherdmania australis sp. nov. Australia, Kott (1) pp. 103-4; E. claviformis larval structure and development, Trason.

Macroclinum arenaceum from N. Auckland, New Zealand, Brewin; M. calypsonis, Pérès (1); M. duboscqi, Pérès (2); M. lacazei sp. nov. E. Mediterranean, Pérès (1) pp. 180-1 fig.; M. seneglense, Pérès (1).

Morchellium argus neural gland, GODEAUX.

Parascidia areolata and P. turbinatum neural gland, Godeaux.

Polyclinum macrophyllum from Mozambique, larva figd., MILLAR (1).

Psammaplidium pedunculatum = Ritterella herdmania nec R. pedunculata, Kott (1).

Pseudodistoma australis sp. nov. Australia, adult and larva, Kott (1) pp. 101-2 fig. 32; P. cyrnusense, Pérès (1, 2).

Ritterella dispar sp. nov. Australia, adult and larva, Kott (1) p. 102 figs. 33-4; R. herdmania nom. nov. for Psammaplidium pedunculatum nec R. pedunculata, Australia, Kott (1).

Sidneioides ivicense sp. nov. E. Mediterranean, Pérès (2) pp. 227-9 fig. 1; S. tamaranae from Australia, larva, Kott (1).

Sigillinaria aurea sp. nov. N. Auckland, New Zealand, larva and adult, Brewin p. 580 fig. 2; S. opaca from N. Auckland, Brewin.

Synoicum hypurgon var. arenosum var. nov. Arabia, Kott (2) p. 141; S. pulmonaria sperm, Franzén.

DIDEMNIDAE

Didemnopsis sansibaricum from Mozambique, Millar (1).

Didemnum candidum from Arabia, Kott (2), from N. Auckland, New Zealand, Brewin, Pérès (1, 2); D. chondrilla from N. Auckland, Brewin; D. dentatum, Pérès (2); D. fulgens, Pérès (1, 2); D. lahillei, Pérès (1); D. maculosum, Pérès (1); D. moseleyi from Arabia, Kott (2); D. paradoxum adult and larva redescribed and figd., Brewin; D. psammatodes from Mozambique, Millar (1), var ianthinum from Arabia, larva figd., Kott (2), var. maculatum from N. Auckland, Brewin.

Diplosoma modestum from Mozambique, larva figd., MILLAR (1); D. spongiforme from Red Sea, Kott (2).

Lissoclinum batailloni, Pérès (2); L. capense from Arabia, Kott (2).

Leptoclinides diemensis from N. Auckland, New Zealand, Brewin.

Leptoclinum as fouling organism, MAWATARI & KOBAYASHI (1, 2).

†Micrascidites gen. nov., De-FLANDRE & DEFLANDRE-RIGAUD p. 48 figs. 1-5, type sp. M. vulgaris; M. vulgaris sp. nov. fossil Pliocene Paris and Australia, spicules, DeFLANDRE & DEFLANDRE-RIGAUD p. 48 figs. 1-5.

Polysyncraton lacazei, Pérès (1, 2).

Trididemnum aspiculatum sp. nov. Arabia: Kuria Muria Islands, Kott (2) pp. 139-40; T. lüderitzi from Arabia, Kott (2); T. tenerum, Pérès (1, 2), sperm, Franzén.

DIAZONIDAE

Rhopalaea neapolitana, Pérès (1, 2).

Rhopalopsis hartmeyeri, Pérès (2).

CIONIDAE

Ciona intestinalis, chemical embryology, Berg, chromosomes, Minganti (2), embryology, Ortolani (2, 4), excretion, Goodbody, heart muscle, electron micrography, Kawaguti & Ikomoto, iodine in endostyle, Barrington, Barrington & Franchi, life cycle, Sabbadin (2), nervous system, Markman, sperm, Franzén; C. savignyi, Pérès (2).

PEROPHORIDAE

Ecteinascidia turbinata ecology, Sourie,

Perophora orientalis heart beat, EBARA (4, 5); P. viridis from Sierra Leone, MILLAR (2).

CORELLIDAE

Corella eumoyta from N. Auckland, New Zealand, Brewin; C. parallelogramma, Pérès (2), sperm, Franzén.

ASCIDIIDAE

Ascidia arenosa from Mozambique, Millar (1); A. aspersa, Pérès (1); A. conchilega, Pérès (2), ecology, Gautier; A. incrassata from Mozambique, Millar (1); A. malaca chromosomes, Minganti (2), from Red Sea, Kott (2); A. melanostoma from Red Sea, Kott (2); A. mentula, Pérès (1, 2), chromosomes, Minganti (2); A. oblique sperm, Franzén; A. virginea, Pérès (2), sperm, Franzén.

Ascidiella aspersa chromosomes, MINGANTI (2), experimental embryology, DALCQ, excretion, GOODBODY, sperm, FRANZÉN; A. pellucida, PÉRÈS (1, 2); A. senegalensis from Sierra Leone, MILLAR (2).

Phallusia mammillata, Pérès (1, 2), chromosomes, Minganti (2), chemical embryology, Mancuso, Vincentiis (2), embryology, Ortolani (1, 3, 4).

BOTRYLLIDAE

Botrylloides nigrum from Red Sea, Kott (2); B. violaceum as a fouling organism, MAWATARI & KOBAYASHI (1, 2). Botryllus communis budding, OKA & WATANABE (3); B. magnicoecus from N. Auckland, New Zealand, BREWIN; B. planus from Mozambique, MILLAR (1); B. primigenus budding, OKA & WATANABE (2, 3), colony specificity, OKA & WATANABE (1); B. schlosseri budding, SABBADIN (1).

Polyandrocarpa inhacae sp. nov. Mozambique, MILLAR (1) pp. 927-9 fig. 14; P. tincta from Mozambique, MILLAR (1).

Symplegma viride from Mozambique, MILLAR (1).

STYELIDAE

Arnbäckia novae-zealandicae from North Auckland, Brewin.

Boltenia echinata sperm, Franzén.

Cnemidocarpa bicornuata, C. madagascariensis var. regalis and C. nisiotis from N. Auckland, New Zealand, Brewin.

Dendroda grossularia embryology, Ivanova-Kasas, sperm, Franzén.

Okamia thilenii from N. Auckland, New Zealand, Brewin.

Polycarpa aurita and P. cryptocarpa from Arabia, Kott (2); P. fibrosa, Pérès (1); P. pomaria, Pérès (1, 2), sperm, Franzén; P. thelypanes from Red Sea, Kott (2); P. translucida from Sierra Leone, Millar (2).

Styela canopus from Red Sea, Kott (2); S. coriacea ecology, Diehl; S. marquesana from Mozambique, Millar (1); S. partita, Pérès (1. 2), sperm. Franzén; S. plicata life cycle Sabbadin (2); S. rustica sperm, Franzén.

Syncarpa longicaudata sp. nov. Kuril Islands, Skalkin pp. 297-9.

Theodorella torus from N. Auckland, New Zealand, Brewin.

PYURIDAE

Halocynthia papillosa, Pérès (1, 2).

Herdmania momus from Mozambique, MILLAR (1), from the Red Sea, Kott (2).

Microcosmus kura from N. Auckland, New Zealand, Brewin; M. sulcatus, Pérès (2).

Pyura cancellata and P. pulla from N. Auckland, New Zealand, Brewin.

P. sansibarica from Mozambique, MILLAR (1); P. subuculata and P. trita from N. Auekland, Brewin.

MOLGULIDAE

Ctenicella appendiculata var. korotneffi, Pérès (1).

Eugyra arenosa sperm, Franzén; E. (Gamaster) peresi sp. nov. Sierra Leone, Millar (2) pp. 374-5 fig. 5.

Molgula citrina sperm, Franzén; M. manhattensis life cycle, Sabbadin (2), excretion, Goodbody, radiation damage, Grosch & Smith; M. mortenseni from N. Auckland, New Zealand, Brewin; M. tubifera from Ostend, Belgium Lefevere et al.

HEXACROBYLIDAE

Hexacrobylus sp. from Arabia, Kott (2).

THALIACEA

PYROSOMIDA

Pyrosoma atlanticum, Trégou-BOFF (1).

DOLIOLIDA

Dolioletta mirabilis from E. Atlantic, Alvariño (2), from W. Mediterranean, Alvariño (1).

Doliolum denticulatum and D. gegenbauri from E. Atlantic, ALVARIÑO; D. nationalis from Mediterranean, TRÉGOUBOFF (1), ALVARIÑO (1), from E. Atlantic, ALVARIÑO (2).

SALPIDA

†Protornis glaronensis shown to be fossil salp of Oligocene, Peyer.

Salpa fusiformis, ALVARIÑO (2), from N. Sea, GLOVER; S. maxima from E. Atlantic, ALVARIÑO (2); S. virgula sperm, Franzén.

LARVACEA

OIKOPLEURIDAE

Haplopleura longicauda from Trinidad, Björnberg & Forneris.

Megalocercus abyssorum from Algiers, Trégouboff.

Oikopleura cophocerca from Trinidad, Björnberg & Forneris; O. dioica from N. Sea, Glover, from Ostend, Belgium, Lefevre et al., from Trinidad, Björnberg & Forneris; O. fusiformis from Trinidad, Björnberg & Forneris; O. labradorensis from N. Sea, Glover; O. parva and O. rufescens from Trinidad, Björnberg & Forneris.

Sinisteroffia gen. nov., Tokioka p. 360, type sp. S. scrippsi; S. scrippsi sp. nov. Peru, Tokioka pp. 360-3 figs. 1-2.

Stegosoma magnum from Trinidad, Björnberg & Forneris.

FRITILLARIIDAE

Fritillaria aequatorialis anatomy, Токлока, from Trinidad, Вубянвенс

& Forneris; F. aplostoma, F. borealis, F. formica, F. megachile, F. messanensis and F. pellucida from Trinidad, Björnberg & Forneris; F. taeniogona sp. nov. Tropical E. Pacifie, Tokioka pp. 363-4 figs. 3-4.

CEPHALOCHORDATA

Amphioxides pelagicus from Brazil, Björnberg (1).

Amphioxus neurosecretion, CEC-CATTY,

Branchiostoma arabiae sp. nov. Arabia and Red Sea, Webb pp. 122-6 fig. 1; B. belcheri redescribed and figd., Webb, chromosomes, Nogusa; B. lanceolatum from Arabia, Webb, sperm, Franzén, structure, Tenbaum.

